

Fracture Liaison Service Database

Commissioners report March 2019

References

1. Kanis JA, Johnell O, De Laet C *et al.* A meta-analysis of previous fracture and subsequent fracture risk. *Bone* 2004;35:375–82.
2. Huntjens KM, Kosar S, van Geel TA *et al.* Risk of subsequent fracture and mortality within 5 years after a non-vertebral fracture. *Osteoporos Int* 2010;21:2075–82.
3. Royal College of Physicians. *National Hip Fracture Database (NHFD) annual report 2017*. London: RCP, 2018.
4. Schnell S, Friedman SM, Mendelson DA, Bingham KW, Kates SL. The 1-year mortality of patients treated in a hip fracture program for elders. *Geriatr Orthop Surg Rehabil* 2010;1:6–14.
5. Keene GS, Parker MJ, Pryor GA. Mortality and morbidity after hip fractures. *BMJ* 1993;307:1248–50.
6. Griffin XL, Parsons N, Achten J, Fernandez M, Costa ML. Recovery of health-related quality of life in a United Kingdom hip fracture population. The Warwick Hip Trauma Evaluation--a prospective cohort study. *Bone Joint J* 2015;97-B:372–82.
7. National Osteoporosis Society. Benefits calculator 2018. Bath: NOS, 2018. <https://benefits.nos.org.uk/> [Accessed 23 January 2019].
8. National Osteoporosis Society. *FLS Pathway and Costing Tool V3.16*. Bath: NOS, 2018.